

Power densities for two-step gamma-ray transitions from isomeric states

Silviu Olariu and Agata Olariu

Institute of Physics and Nuclear Engineering,

Department of Fundamental Experimental Physics

76900 Magurele, P.O. Box MG-6, Bucharest, Romania

e-mail: olariu@ifin.nipne.ro

Abstract

We have calculated the incident photon power density P_2 for which the two-step induced emission rate from an isomeric nucleus becomes equal to the natural isomeric decay rate. We have analyzed two-step transitions for isomeric nuclei with a half-life greater than 10 min, for which there is an intermediate state of known energy, spin and half-life, for which the intermediate state is connected by a known gamma-ray transition to the isomeric state and to at least another intermediate state, and for which the relative intensities of the transitions to lower states are known. For the isomeric nucleus ^{166m}Ho , which has a 1200 y isomeric state at 5.98 keV, we have found a value of $P_2 = 6.3 \times 10^7 \text{ W cm}^{-2}$, the intermediate state being the 263.8 keV level. We have found power densities P_2 of the order of $10^{10} \text{ W cm}^{-2}$ for several other isomeric nuclei.

PACS numbers: 23.20.Lv, 23.20.Nx, 25.20.Dc, 42.55.Vc

The induced deexcitation of an isomeric nucleus is a two-step process in which the nucleus, initially in the isomeric state $|i\rangle$, first absorbs a photon of energy E_{ni} to reach a higher intermediate state $|n\rangle$, then the nucleus makes a transition to a lower state $|l\rangle$ by the emission of a gamma-ray photon of energy E_{nl} or by internal conversion. In a previous work [1] it has been required that the cascade originating on the state $|n\rangle$ should contain a gamma-ray transition of energy $E_\gamma > 2E_{ni}$, the latter relation representing a condition of upconversion of the energy of the incident photons, and it was found that in favorable cases the two-step induced emission rates become equal to the natural isomeric decay rates for incident power densities of the order of 10^{10} W cm $^{-2}$. The Weisskopf estimates were used in [1] in cases when tabulated half-lives or partial widths were not available.

In this report we have analyzed, regardless of the upconversion condition $E_\gamma > 2E_{ni}$, the two-step transitions for isomeric nuclei with a half-life greater than 10 min, for which there is an intermediate state $|n\rangle$ of known energy, spin and half-life, for which the intermediate state $|n\rangle$ is connected by a known gamma-ray transition to the isomeric state and to a lower state $|l\rangle$, and for which the relative intensities of the transitions from the intermediate state $|n\rangle$ are known. [2] We have calculated the incident power density P_2 for which the two-step induced emission rate becomes equal to the natural isomeric decay rate. We have found a value of $P_2 = 6.3 \times 10^7$ W cm $^{-2}$ for ^{166}Ho , which has a 1200 y isomeric state at $E_i=5.98$ keV, and the intermediate state is the $E_n=263.8$ keV level, so that $E_{ni} = 257.8$ keV. We have also found several isomers for which the power density P_2 is of the order of 10^{10} W cm $^{-2}$.

We assume that the nucleus is initially in an isomeric state $|i\rangle$ of energy E_i , spin J_i and half-life t_i . By absorbing an incident photon of energy E_{ni} , the nucleus makes a transition to a higher intermediate state $|n\rangle$ of energy E_n , spin J_n and half-life t_n . The state $|n\rangle$ then decays into a lower state $|l\rangle$ of energy E_l and spin J_l by the emission of a gamma-ray photon having the energy E_{nl} or by internal conversion, as shown in Fig. 1. In some cases the state $|l\rangle$ may be situated above the isomeric state $|i\rangle$, and in these cases the transition $|n\rangle \rightarrow |l\rangle$ is followed by further gamma-ray transitions to lower states.

As shown in [1], the rate $w_{il}^{(2)}$ for the two-step transition $|i\rangle \rightarrow |n\rangle \rightarrow |l\rangle$ is

$$w_{il}^{(2)} = \sigma_{ni} \hbar \Gamma_{eff} N(E_{ni}). \quad (1)$$

where

$$\sigma_{ni} = \frac{2J_n + 1}{2J_i + 1} \frac{\pi^2 c^2 \hbar^2}{E_{ni}^2} \quad (2)$$

is the induced-emission cross section for the transition $|i\rangle \rightarrow |n\rangle$. The quantity Γ_{eff} is the effective width of the two-step transition,

$$\Gamma_{eff} = F_R \ln 2/t_n, \quad (3)$$

where the dimensionless quantity F_R has the expression

$$F_R = \frac{(1 + \alpha_{nl})R_{ni}R_{nl}}{[(1 + \alpha_{ni})R_{ni} + (1 + \alpha_{nl})R_{nl} + \sum_{l'}(1 + \alpha_{nl'})R_{nl'}]^2}, \quad (4)$$

$R_{ni}, R_{nl}, R_{nl'}$ being the relative gamma-ray intensities and $\alpha_{ni}, \alpha_{nl}, \alpha_{nl'}$ the internal conversion coefficients for the transitions $|n\rangle \rightarrow |i\rangle, |n\rangle \rightarrow |l\rangle, |n\rangle \rightarrow |l'\rangle, l' \neq i, l$. In Eq. (1) $N(E_{ni})$ is the spectral intensity for the incident photon flux, defined such that $N(E)dE$ should represent the number of photons incident per unit surface and time and having the energy between E and $E + dE$.

The spectral intensity N_2 for which the two-step transition rate $w_{il}^{(2)}$ becomes equal to the natural decay rate $\ln 2/t_i$ of the isomeric nucleus is

$$N_2 = \frac{\ln 2}{\sigma_{int} t_i}, \quad (5)$$

where the integrated cross section σ_{int} is

$$\sigma_{int} = \sigma_{ni} \hbar \Gamma_{eff}. \quad (6)$$

The incident power density P_2 for which the induced emission rate becomes equal to the natural decay rate of the isomeric state can then be estimated as

$$P_2 = N_2(E_{ni}) E_{ni}^2. \quad (7)$$

We have analyzed two-step transitions for isomeric nuclei with a half-life greater than 10 min, for which there is an intermediate state $|n\rangle$ of known energy E_n , spin J_n and half-life t_n , for which the intermediate state is connected by a known gamma-ray transition to the isomeric state $|i\rangle$ and to a lower state $|l\rangle$, and for which the relative intensities $R_{ni}, R_{nl}, R_{nl'}$ of the transitions to lower states are known. The internal conversion coefficients have been calculated by interpolation from refs. [3] and [4]. The cases for which all the previously mentioned quantities were available in

ref. [2] are listed in Table I. If the input values were available for several two-step transitions of a certain isomeric nucleus, we gave the two-step transition having the lowest value of P_2 . The two-step transitions for the nuclei ^{52}Mn , ^{99}Tc , ^{152}Eu , ^{178}Hf , ^{201}Bi , ^{204}Pb , which fulfil the upconversion condition, have been studied in ref. [1] and are not included in Table I. If the half-life t_n of the intermediate state was given in [2] as less than ($<$) or greater than ($>$) a certain value, we have calculated $\hbar\Gamma_{eff}$, σ_{int} and P_2 corresponding to the given limiting value of t_n . In the case of ^{166}Ho we have neglected the 3.1 keV transition, of unknown intensity, from the 263.8 keV level. In the case of ^{97}Tc the intensity of the 441.2 keV transition from the 656.9 keV level was given as an upper limit. In the case of ^{121}Sn we have neglected the weak 56.35 keV transition from the 925.6 keV level. The half-life of the 0.0768 keV isomeric level of ^{235}U and the intensity of the 637.7 keV transition from the 637.79 keV intermediate state are approximate values. In the case of ^{34}Cl the intensity of the 2433.8 keV transition from the 2580.2 keV intermediate state was given as an upper limit, as are the intensities of some other transitions from this intermediate state. In the case of ^{123}Sn we have neglected the weak 284.7 keV transition from the 1130.5 keV intermediate state.

For the isomeric nucleus ^{166m}Ho , which has a 1200 y isomeric state at $E_i=5.98$ keV, we have found a value of $P_2 = 6.3 \times 10^7 \text{ W cm}^{-2}$, the intermediate state being the $E_n=263.8$ keV level. The relatively low value of P_2 is due to the long half-life of this isomer. The largest effective width in Table I is for ^{34m}Cl , for which $\hbar\Gamma_{eff} = 2.5 \times 10^{-3} \text{ eV}$. The relatively large effective width is due to the short half-life of the intermediate state, $t_n=3$ fs. The energy of the pumping transition is however large, $E_{ni}=2433.8$ keV, and the multipolarity is E2/M1. The large value of P_2 for ^{34}Cl is due to the relatively small half-life of the isomeric state. There are several cases in Table I for which the power density P_2 is of the order of $10^{10} \text{ W cm}^{-2}$. For ^{97}Tc and ^{95}Tc the low values of P_2 can be attributed to the short half-lives of the intermediate state, whereas for ^{113}Cd and ^{121}Sn the values of P_2 are a result of the long isomeric half-lives. The two-step transitions for which all the input values were available represent a small fraction of the total number of possible two-step transitions in isomeric nuclei.

Since the number of isomeric nuclei in a sample is limited by the total activity of that sample, a longer isomeric half-life means a larger number of isomeric nuclei in the sample. The two-step induced transitions can be induced by incident photons, as assumed in this work, or directly by


incident electrons. The cross sections for the two-step electron excitation of isomeric states are about two orders of magnitudes lower than the cross sections for two-step photon excitation but the two-step photon excitation rates are also proportional to the efficiency with which the energy of an incident electron beam can be converted into bremsstrahlung.

References

- [1] S. Olariu and A. Olariu, Phys. Rev. C **57**, accepted for publication (1998).
- [2] *Table of Isotopes*, 8th ed., edited by R. Firestone (Wiley, New York, 1996).
- [3] I. M. Band, M. B. Trzhaskovskaya and M. A. Listengarten, Atomic Data and Nuclear Data Tables **18**, 433 (1976).
- [4] F. Rösels, H. M. Fries, K. Alder and H. C. Pauli, Atomic Data and Nuclear Data Tables **21**, 91 (1978), **21**, 291 (1978).

Fig. 1. Two-step deexcitation of an isomeric nucleus. The nucleus, initially in the isomeric state $|i\rangle$, makes a transition to a higher nuclear state $|n\rangle$ by the absorption of an incident photon of energy E_{ni} , then it emits a photon of energy E_{nl} to reach the lower nuclear state $|l\rangle$.

Table I. Power density P_2 for which the two-step induced emission rate becomes equal to the natural decay rate of the isomeric state of energy E_i and half-life t_i . The intermediate state of the problem is the level of energy E_n and half-life t_n . The multipolarities of the transitions of energy E_{ni}, E_{nl} are given in the multipole column.



Nucleus	E_i (keV)	t_i (s)	E_n (keV)	t_n (s)	E_{ni} (keV)	E_{nl} (keV)	multipole ni/nl	F_R	$\hbar\Gamma_{eff}$ (eV)	σ_{int} (cm ² eV)	P_2 (W cm ⁻²)
¹⁶⁶ Ho	6.0	3.8×10 ¹⁰	263.8	<5.0×10 ⁻¹⁰	257.8	72.9	M2/E2	8.0×10 ⁻²	7.3×10 ⁻⁸	3.1×10 ⁻²⁷	6.3×10 ⁷
⁹⁷ Tc	96.6	7.8×10 ⁶	656.9	>7.6×10 ⁻¹³	560.3	441.2	E2/E1	2.8×10 ⁻²	1.7×10 ⁻⁵	6.1×10 ⁻²⁵	7.3×10 ⁹
⁹⁵ Tc	38.9	5.3×10 ⁶	927.8	>5.9×10 ⁻¹³	888.9	301.1	E1/M1	1.4×10 ⁻¹	1.1×10 ⁻⁴	1.0×10 ⁻²⁴	1.6×10 ¹⁰
¹¹³ Cd	263.6	4.4×10 ⁸	522.3	3.2×10 ⁻¹⁰	258.8	206.4	E2/E1	1.1×10 ⁻²	1.6×10 ⁻⁸	5.9×10 ⁻²⁸	2.8×10 ¹⁰
¹²¹ Sn	6.3	1.7×10 ⁹	925.6	2.5×10 ⁻¹⁰	919.3	925.6	M2/E2	3.7×10 ⁻²	6.7×10 ⁻⁸	2.0×10 ⁻²⁸	2.7×10 ¹¹
⁵⁸ Co	24.9	3.3×10 ⁴	373.9	6.0×10 ⁻¹³	349.1	320.8	M1/M1	3.2×10 ⁻²	2.4×10 ⁻⁵	7.6×10 ⁻²⁵	5.4×10 ¹¹
⁸³ Kr	41.5	6.6×10 ³	562.0	6.0×10 ⁻¹²	520.4	552.6	E2/E1	1.9×10 ⁻¹	1.5×10 ⁻⁵	6.2×10 ⁻²⁵	7.4×10 ¹²
¹³¹ Xe	163.9	1.0×10 ⁶	666.9	<5.0×10 ⁻¹⁰	503.0	325.8	E2/M1	2.4×10 ⁻¹	2.2×10 ⁻⁷	2.2×10 ⁻²⁷	1.2×10 ¹³
¹⁸⁹ Os	30.8	2.1×10 ⁴	219.4	1.9×10 ⁻¹⁰	188.6	219.4	M1/E2	4.1×10 ⁻²	9.9×10 ⁻⁸	8.6×10 ⁻²⁷	2.2×10 ¹³
¹¹⁹ Sn	89.5	2.5×10 ⁷	787.0	1.9×10 ⁻¹⁰	697.4	763.1	M2/E2	4.8×10 ⁻³	1.2×10 ⁻⁸	6.1×10 ⁻²⁹	3.5×10 ¹³
²³⁵ U	0.0768	1.5×10 ³	637.79	1.3×10 ⁻¹¹	637.72	637.79	E1/E2	1.2×10 ⁻¹	4.1×10 ⁻⁶	7.8×10 ⁻²⁶	3.9×10 ¹⁴
³⁴ Cl	146.4	1.9×10 ³	2580.3	<3×10 ⁻¹⁵	2433.8	2580.2	E2/M1	1.7×10 ⁻²	2.5×10 ⁻³	7.0×10 ⁻²⁵	4.9×10 ¹⁴
¹²⁵ Te	144.8	5.0×10 ⁶	642.2	<6.0×10 ⁻¹⁰	497.4	606.7	M2/E2	1.3×10 ⁻³	9.7×10 ⁻¹⁰	1.0×10 ⁻²⁹	5.5×10 ¹⁴
¹¹⁷ Sn	314.6	1.2×10 ⁶	711.5	9.8×10 ⁻¹⁰	396.6	553.0	M2/E2	1.7×10 ⁻³	8.0×10 ⁻¹⁰	1.3×10 ⁻²⁹	1.1×10 ¹⁵
⁸⁷ Sr	388.5	1.0×10 ⁴	2414.5	1.3×10 ⁻¹³	2026.0	643.8	M1/E1	5.7×10 ⁻³	2.0×10 ⁻⁵	3.8×10 ⁻²⁶	1.2×10 ¹⁵
¹²⁴ Sb	36.8	1.2×10 ³	125.2	8.6×10 ⁻⁸	88.4	37.6	E2/E2	5.1×10 ⁻²	2.7×10 ⁻¹⁰	1.0×10 ⁻²⁸	7.0×10 ¹⁵
¹²³ Sn	24.6	2.4×10 ³	1155.0	<1.0×10 ⁻¹⁰	1130.5	536.4	E2/E1	1.4×10 ⁻²	6.2×10 ⁻⁸	3.7×10 ⁻²⁸	1.6×10 ¹⁷
¹⁰³ Rh	39.8	3.4×10 ³	650.1	<1.0×10 ⁻¹⁰	610.3	292.7	M1/E1	3.8×10 ⁻⁴	1.7×10 ⁻⁹	1.3×10 ⁻²⁹	9.2×10 ¹⁷
¹¹⁷ In	315.3	7.0×10 ³	1891.9	4.2×10 ⁻¹⁰	1576.6	840.2	E1/E2	8.6×10 ⁻³	9.4×10 ⁻⁹	1.4×10 ⁻²⁹	2.7×10 ¹⁸